



TS-205

Black Eye™

Endoscopic **Marker**

Indication for use

Black Eye is indicated for endoscopically marking lesions in the GI tract permanently.

Black Eye's biocompatible, non-toxic formulation contains no harmful ingredient. Black Eye consists of medical grade carbon black which lasts for more than 10 years in the body!

Permanent Ink

Stable Suspension

Poor suspension can make the solution separated, which leads to unstable marking!
"No shake" Necessary = Easier, Faster, Safer!

Poor emulsion can choke the tube of the syringe or body fluid!

Stable Emulsion

Proven in over 50 countries



Black Eye™

Endoscopic Marker



STORAGE

Store the product at room temperature.

SHELF LIFE

24 months.

PHYSICAL PROPERTIES

- 1) Color: Black
- 2) Density: 1.05g/mL
- 3) Boiling Point: 97°C (207°F)
- 4) Freezing Point: -0.5°C (33°F)

STERILIZATION

- 1) Steam sterilization
- 2) Sterility Assurance Level (SAL): 10^{-6}

RAW MATERIALS

Water for injection
Glycerin
Carbon Black

DIRECTIONS FOR USE

A 23-25 gauge sclerotherapy needle with a needle length of 4mm or less is recommended for this procedure.

IMPORTANT

When the needle is properly positioned, insert the tip 30°-45° tangentially into the submucosal space of the colon.

CAUTION

Inject 0.5~0.75ml of the marker into each injection site. This volume typically produces a mark on the colon measuring 1~2 cm in diameter. The maximum recommended dose is 8ml per patient.



5ml syringe, box of 10

Rectangular canister for secure and compact storage. Blister packaging acts as a sterile barrier and protects against cross-contamination.



THE STANDARD CO., LTD.

120, Gunpocheomdansaneop 2-ro, Gunpo-si, Gyeonggi-do, 15880, Korea
Tel : +82-2-838-5533 Fax : +82-2-838-5523 E-mail : sales1@thestd.co.kr